



REFERENCE DECK

1. Antonio G. Richetta¹, Carmen Cantisani¹, *, Vincent W. Li², Carlo Mattozzi¹, Luca Melis¹, Federico De Gado¹, Simona Giancristoforo¹, Emidio Silvestri¹ and Stefano Calvieri¹ (2011), “Hydrofiber Dressing andWound Repair: Review of the Literature and New Patents”, 1Rome, Italy, 2 Boston, MA, USA, DOI: <https://doi.org/10.2174/187221311795399264>.
2. Alison Freeman, Community Staff Nurse, Walsall Healthcare NHS Trust., Clare Morris, Senior Clinical Services Manager, (2018), “A clinical evaluation of a hydroactive fibre dressing in the management of lowerlimb chronic wounds in a primary care setting”, Harrogate, UK, Harrogate, UK.
3. Carolina D. Weller^{1*}, Victoria Team¹ and Geoffrey Sussman² (2020), “First-Line Interactive Wound Dressing Update: A Comprehensive Review of the Evidence”, Melbourne, VIC, Australia, Front. Pharmacol., DOI: <https://doi.org/10.3389/fphar.2020.00155>.
4. Julian F. Guest and Francis J. Ruiz (2005), “Modelling the cost implications of using carboxymethylcellulosedressing compared with gauze in the management of surgical wounds healing by secondary intention in the US and UK”, Northwood, Middlesex, UK, DOI: <https://doi.org/10.1185/030079905X25532>.
5. K. Reddersen¹, C. Wiegand¹, S. Springer¹, M. Abel², P. Ruth², U.-Ch. Hipler¹(2014), “In-vitro-analysis ofthe fluid management by hydroactive wound dressings using a maceration model”, Madrid/Spain, 24th Conference of the European Wound Management Association (EWMA), 14.-16. May 2014.
6. Raman Mundi¹, Harman Chaudhry¹, Seper Ekhtiari², Prabjit Ajrawat¹, Daniel M Tushinski², Thomas J Wood² and Mohit Bhandari² (2021), “Efficacy of hydrofibre dressing following total joint arthroplasty: ameta-analysis of randomised controlled trials”, Toronto, ON, Canada, DOI: <https://doi.org/10.1177/11207000211012669>.
7. Reiva Farah Dwiyana, 1; Yuri Yogyo, 1; *Srie Prihanti Gondokaryono, 1; Inne Arline Diana, 1; Oki Suwarsa,1; Lies Marlysa Ramali, 1; Eva Krishna Sutedja, 1; July Iriani Rahardja,1; Hendra Gunawan, 1 (2019), “Clinical efficacy of biocellulose, carboxymethyl cellulose and normal saline dressing in epidermolysis bullosa”, Bandung, Indonesia, 194.066.072.104, journal of wound care silk road supplement vol 28, no 10.
8. Stefanie de Lange¹, Michael Schmitz², Martin Abel¹(2017), “Assessment of efficacy and tolerability of anew developed hydroactive wound dressing”, 1 Lohmann & Rauscher, Rengsdorf, Deutschland; 2 MCS Medical Consulting, Oberahr, Deutschland, Journée Cicatrisations (CICA) Paris, 15 -17 January 2017.
9. Tin Wui Wong Nor Amlizan Ramli (2014), “Carboxymethylcellulose film for bacterial wound infectioncontrol and healing”, Selangor, Malaysia, DOI: <http://dx.doi.org/doi:10.1016/j.carbpol.2014.06.002>.
10. Vimala Kanikireddy a, *, Kokkarachedu Varaprasad b, *, Tippabattini Jayaramudu c, Chandrasekaran Karthikeyan b, Rotimi Sadiku d (2020), “Carboxymethyl cellulose-based materials for infection control andwound healing: A review”, a Telangana, India, b Avenida Collao 1202, Concepción, Chile c deTalca, 747, Talca, Chile d Tshwane University of Technology, Pretoria West Campus, Staatsarillerie Rd, Pretoria 1083, South Africa, DOI: <https://doi.org/10.1016/j.ijbiomac.2020.07.160>.



REFERENCE DECK

11. Wolber Silke 1, Scheuenstuhl Yvonne 2, de Lange Stefanie 3, Abel Martin 3(2017), “Case series demonstrating the performance and safety of a novel superabsorbing dressing* in high exuding chronic wounds – first results”, 1 Diabetologie Wetbergen, Hannover; 2 Praxis Dres. Römer/Michel, Zollingen, 3 Medical & Regulatory Affairs, Lohmann & Rauscher GmbH & Co. KG, Rengsdorf, Deutschland, Journées Cicatrisation, Paris, 15-17 January 2017.
12. Yoonjin Kim, Song Jun Doh, Gyu Dong Lee, Chaehwa Kim, and Jung Nam Im*, (2019), “Composite Nonwovens Based on Carboxymethyl Cellulose for Wound Dressing Materials”, Ansan 15588, Korea, DOI:<https://doi.org/10.1007/s12221-019-9261-9>.